



Some topological properties of fuzzy cone metric spaces

Tarkan Öner

Department of Mathematics, Faculty of Sciences, Muğla Sıtkı Koçman University, 48000 Mugla, Turkey.

Communicated by R. Saadati

Abstract

We prove Baire's theorem for fuzzy cone metric spaces in the sense of Öner *et al.* [T. Öner, M. B. Kandemir, B. Tanay, J. Nonlinear Sci. Appl., **8** (2015), 610–616]. A necessary and sufficient condition for a fuzzy cone metric space to be precompact is given. We also show that every separable fuzzy cone metric space is second countable and that a subspace of a separable fuzzy cone metric space is separable. ©2016 All rights reserved.

Keywords: Fuzzy cone metric space, Baire's theorem, separable, second countable.
2010 MSC: 54A40, 54E35, 54E15, 54H25.

1. Introduction

After Zadeh [13] introduced the theory of fuzzy sets, many authors have introduced and studied several notions of metric fuzziness ([2], [3], [4], [8], [9]) and metric cone fuzziness ([1], [10] from different points of view).

By modifying the concept of metric fuzziness introduced by George and Veeramani [4], Öner *et al.* [10] studied the notion of fuzzy cone metric spaces. In particular, they proved that every fuzzy cone metric space generates a Hausdorff first-countable topology.

Here we study further topological properties of these spaces whose fuzzy metric version can be found in [4], [5] and [6]. We show that every closed ball is a closed set and prove Baire's theorem for fuzzy cone metric spaces. Moreover, we prove that a fuzzy cone metric space is precompact if and only if every sequence in it has a Cauchy subsequence. Further, we show that $X_1 \times X_2$ is a complete fuzzy cone metric space if and only if X_1 and X_2 are complete fuzzy cone metric spaces. Finally it is proven that every separable fuzzy cone metric space is second countable and a subspace of a separable fuzzy cone metric space is separable.

Email address: tarkanoner@mu.edu.tr (Tarkan Öner)

2. Preliminaries

Let E be a real Banach space, θ the zero of E and P a subset of E . Then P is called a cone [7] if and only if

- 1) P is closed, nonempty, and $P \neq \{\theta\}$;
- 2) if $a, b \in \mathbb{R}$, $a, b \geq 0$ and $x, y \in P$, then $ax + by \in P$;
- 3) if both $x \in P$ and $-x \in P$, then $x = \theta$.

Given a cone P , a partial ordering \preceq on E with respect to P is defined by $x \preceq y$ if only if $y - x \in P$. The notation $x \prec y$ will stand for $x \preceq y$ and $x \neq y$, while $x \ll y$ will stand for $y - x \in \text{int}(P)$ [7]. Throughout this paper, we assume that all the cones have nonempty interiors.

There are two kinds of cones: normal and nonnormal ones. A cone P is called normal if there exists a constant $K \geq 1$ such that for all $t, s \in E$, $\theta \preceq t \preceq s$ implies $\|t\| \leq K\|s\|$, and the least positive number K having this property is called normal constant of P [7]. It is clear that $K \geq 1$ [11].

According to [12], a binary operation $* : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is a continuous t -norm if it satisfies:

- 1) $*$ is associative and commutative;
- 2) $*$ is continuous;
- 3) $a * 1 = a$ for all $a \in [0, 1]$;
- 4) $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$, $a, b, c, d \in [0, 1]$.

In [10], we generalized the concept of fuzzy metric space of George and Veeramani by replacing the $(0, \infty)$ interval by $\text{int}(P)$ where P is a cone as follows:

A fuzzy cone metric space is a 3-tuple $(X, M, *)$ such that P is a cone of E , X is nonempty set, $*$ is a continuous t -norm and M is a fuzzy set on $X^2 \times \text{int}(P)$ satisfying the following conditions, for all $x, y, z \in X$ and $t, s \in \text{int}(P)$ (that is $t \gg \theta$, $s \gg \theta$)

- FCM1) $M(x, y, t) > 0$;
- FCM2) $M(x, y, t) = 1$ if and only if $x = y$;
- FCM3) $M(x, y, t) = M(y, x, t)$;
- FCM4) $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$;
- FCM5) $M(x, y, \cdot) : \text{int}(P) \rightarrow [0, 1]$ is continuous.

If $(X, M, *)$ is a fuzzy cone metric space, we will say that M is a fuzzy cone metric on X .

In [10], it was proven that every fuzzy cone metric space $(X, M, *)$ induces a Hausdorff first-countable topology τ_{fc} on X which has as a base the family of sets of the form $\{B(x, r, t) : x \in X, 0 < r < 1, t \gg \theta\}$, where $B(x, r, t) = \{y \in X : M(x, y, t) > 1 - r\}$ for every r with $0 < r < 1$ and $t \gg \theta$.

A fuzzy cone metric space $(X, M, *)$ is called complete if every Cauchy sequence in it is convergent, where a sequence $\{x_n\}$ is said to be a Cauchy sequence if for any $\varepsilon \in (0, 1)$ and any $t \gg \theta$ there exists a natural number n_0 such that $M(x_n, x_m, t) > 1 - \varepsilon$ for all $n, m \geq n_0$, and a sequence $\{x_n\}$ is said to converge to x if for any $t \gg \theta$ and any $r \in (0, 1)$ there exists a natural number n_0 such that $M(x_n, x, t) > 1 - r$ for all $n \geq n_0$ [10].

A sequence $\{x_n\}$ converges to x if and only if $\lim_{n \rightarrow \infty} M(x_n, x, t) \rightarrow 1$ for each $t \gg \theta$ [10].

3. Results

Definition 3.1. Let $(X, M, *)$ be a fuzzy cone metric space. For $t \gg \theta$, the closed ball $B[x, r, t]$ with center x and radius $r \in (0, 1)$ is defined by $B[x, r, t] = \{y \in X : M(x, y, t) \geq 1 - r\}$.

Lemma 3.2. Every closed ball in a fuzzy cone metric space $(X, M, *)$ is a closed set.

Proof. Let $y \in \overline{B[x, r, t]}$. Since X is first countable, there exists a sequence $\{y_n\}$ in $B[x, r, t]$ converging to y . Therefore $M(y_n, y, t)$ converges to 1 for all $t \gg \theta$. For a given $\epsilon \gg 0$, we have

$$M(x, y, t + \epsilon) \geq M(x, y_n, t) * M(y_n, y, \epsilon).$$

Hence

$$M(x, y, t + \epsilon) \geq \lim_{n \rightarrow \infty} M(x, y_n, t) * \lim_{n \rightarrow \infty} M(y_n, y, \epsilon) \geq (1 - r) * 1 = 1 - r.$$

(If $M(x, y_n, t)$ is bounded, then the sequence $\{y_n\}$ has a subsequence, which we again denote by $\{y_n\}$, for which $\lim_{n \rightarrow \infty} M(x, y_n, t)$ exists.) In particular for $n \in \mathbb{N}$, take $\epsilon = \frac{t}{n}$. Then

$$M(x, y, t + \frac{t}{n}) \geq (1 - r).$$

Hence

$$M(x, y, t) \geq \lim_{n \rightarrow \infty} M(x, y, t + \frac{t}{n}) \geq 1 - r.$$

Thus $y \in B[x, r, t]$. Therefore $B[x, r, t]$ is a closed set. □

Theorem 3.3 (Baire’s theorem). *Let $(X, M, *)$ be a complete fuzzy cone metric space. Then the intersection of a countable number of dense open sets is dense.*

Proof. Let X be the given complete fuzzy cone metric space, B_0 a nonempty open set, and D_1, D_2, D_3, \dots dense open sets in X . Since D_1 is dense in X , we have $B_0 \cap D_1 \neq \emptyset$. Let $x_1 \in B_0 \cap D_1$. Since $B_0 \cap D_1$ is open, there exist $0 < r_1 < 1, t_1 \gg \theta$ such that $B(x_1, r_1, t_1) \subset B_0 \cap D_1$. Choose $r'_1 < r_1$ and $t'_1 = \min\{t_1, t_1/\|t_1\|\}$ such that $B[x_1, r'_1, t'_1] \subset B_0 \cap D_1$. Let $B_1 = B(x_1, r'_1, t'_1)$. Since D_2 is dense in X , we have $B_1 \cap D_2 \neq \emptyset$. Let $x_2 \in B_1 \cap D_2$. Since $B_1 \cap D_2$ is open, there exist $0 < r_2 < 1/2$ and $t_2 \gg \theta$ such that $B(x_2, r_2, t_2) \subset B_1 \cap D_2$. Choose $r'_2 < r_2$ and $t'_2 = \min\{t_2, t_2/2\|t_2\|\}$ such that $B[x_2, r'_2, t'_2] \subset B_1 \cap D_2$. Let $B_2 = B(x_2, r'_2, t'_2)$. Similarly proceeding by induction, we can find an $x_n \in B_{n-1} \cap D_n$. Since $B_{n-1} \cap D_n$ is open, there exist $0 < r_n < 1/n, t_n \gg \theta$ such that $B(x_n, r_n, t_n) \subset B_{n-1} \cap D_n$. Choose an $r'_n < r_n$ and $t'_n = \min\{t_n, t_n/n\|t_n\|\}$ such that $B[x_n, r'_n, t'_n] \subset B_{n-1} \cap D_n$. Let $B_n = B(x_n, r'_n, t'_n)$. Now we claim that $\{x_n\}$ is a Cauchy sequence. For a given $t \gg \theta, 0 < \epsilon < 1$, choose an n_0 such that $t/n_0\|t\| \ll t, 1/n_0 < \epsilon$. Then for $n \geq n_0, m \geq n$, we have

$$M(x_n, x_m, t) \geq M\left(x_n, x_m, \frac{t}{n_0\|t\|}\right) \geq 1 - \frac{1}{n} \geq 1 - \epsilon.$$

Therefore $\{x_n\}$ is a Cauchy sequence. Since X is complete, $x_n \rightarrow x$ in X . But $x_k \in B[x_n, r'_n, t'_n]$ for all $k \geq n$. Since $B[x_n, r'_n, t'_n]$ is closed, $x \in B[x_n, r'_n, t'_n] \subset B_{n-1} \cap D_n$ for all n . Therefore $B_0 \cap (\bigcap_{n=1}^\infty D_n) \neq \emptyset$. Hence $\bigcap_{n=1}^\infty D_n$ is dense in X . □

Definition 3.4. A fuzzy cone metric space $(X, M, *)$ is called precompact if for each r , with $0 < r < 1$, and each $t \gg \theta$, there is a finite subset A of X , such that $X = \bigcup_{a \in A} B(a, r, t)$. In this case, we say that M is a precompact fuzzy cone metric on X .

Lemma 3.5. *A fuzzy cone metric space is precompact if and only if every sequence has a Cauchy subsequence.*

Proof. Suppose that $(X, M, *)$ is a precompact fuzzy cone metric space. Let $\{x_n\}$ be a sequence in X . For each $m \in \mathbb{N}$ there is a finite subset A_m of X such that $X = \bigcup_{a \in A_m} B(a, 1/m, t_0/m\|t_0\|)$ where $t_0 \gg \theta$ is a constant. Hence, for $m = 1$, there exists an $a_1 \in A_1$ and a subsequence $\{x_{1(n)}\}$ of $\{x_n\}$ such that $x_{1(n)} \in B(a_1, 1, t_0/\|t_0\|)$ for every $n \in \mathbb{N}$. Similarly, there exist an $a_2 \in A_2$ and a subsequence $\{x_{2(n)}\}$ of $\{x_{1(n)}\}$ such that $x_{2(n)} \in B(a_2, 1/2, t_0/2\|t_0\|)$ for every $n \in \mathbb{N}$. By continuing this process, we get that for $m \in \mathbb{N}, m > 1$, there is an $a_m \in A_m$ and a subsequence $\{x_{m(n)}\}$ of $\{x_{m-1(n)}\}$ such that $x_{m(n)} \in B(a_m, 1/m, t_0/m\|t_0\|)$ for every $n \in \mathbb{N}$. Now, consider the subsequence $\{x_{n(n)}\}$ of $\{x_n\}$. Given r with $0 < r < 1$ and $t \gg \theta$ there is an $n_0 \in \mathbb{N}$ such that $(1 - 1/n_0) * (1 - 1/n_0) > 1 - r$ and $2t_0/n_0\|t_0\| \ll t$. Then, for every $k, m \geq n_0$, we have

$$\begin{aligned}
 M(x_{k(k)}, x_{m(m)}, t) &\geq M\left(x_{k(k)}, x_{m(m)}, \frac{2t_0}{n_0 \|t_0\|}\right) \\
 &\geq M\left(x_{k(k)}, a_{n_0}, \frac{t_0}{n_0 \|t_0\|}\right) * M\left(a_{n_0}, x_{m(m)}, \frac{t_0}{n_0 \|t_0\|}\right) \\
 &\geq \left(1 - \frac{1}{n_0}\right) * \left(1 - \frac{1}{n_0}\right) \\
 &> 1 - r.
 \end{aligned}$$

Hence $(x_{n(n)})$ is a Cauchy sequence in $(X, M, *)$.

Conversely, suppose that $(X, M, *)$ is a nonprecompact fuzzy cone metric space. Then there exist an r with $0 < r < 1$ and $t \gg \theta$ such that for each finite subset A of X , we have $X \neq \bigcup_{a \in A} B(a, r, t)$. Fix $x_1 \in X$. There is an $x_2 \in X - B(x_1, r, t)$. Moreover, there is an $x_3 \in X - \bigcup_{k=1}^2 B(x_k, r, t)$. By continuing this process, we construct a sequence $\{x_n\}$ of distinct points in X such that $x_{n+1} \notin \bigcup_{k=1}^n B(x_k, r, t)$ for every $n \in \mathbb{N}$. Therefore $\{x_n\}$ has no Cauchy subsequence. This completes the proof. \square

Lemma 3.6. *Let $(X, M, *)$ be a fuzzy cone metric space. If a Cauchy sequence clusters around a point $x \in X$, then the sequence converges to x .*

Proof. Let $\{x_n\}$ be a Cauchy sequence in $(X, M, *)$ having a cluster point $x \in X$. Then, there is a subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ that converges to x with respect to τ_{fc} . Thus, given r with $0 < r < 1$ and $t \gg \theta$, there is an $n_0 \in \mathbb{N}$ such that for each $n \geq n_0$, $M(x, x_{k(n)}, t/2) > 1 - s$ where $s > 0$ satisfies $(1 - s) * (1 - s) > 1 - r$. On the other hand, there is $n_1 \geq k(n_0)$ such that for each $n, m \geq n_1$, we have $M(x_n, x_m, t/2) > 1 - s$. Therefore, for each $n \geq n_1$, we have

$$\begin{aligned}
 M(x, x_n, t) &\geq M(x, x_{k(n)}, \frac{t}{2}) * M(x_{k(n)}, x_n, \frac{t}{2}) \\
 &\geq (1 - s) * (1 - s) \\
 &> 1 - r.
 \end{aligned}$$

We conclude that the Cauchy sequence $\{x_n\}$ converges to x . \square

Proposition 3.7. *Let $(X_1, M_1, *)$ and $(X_2, M_2, *)$ be fuzzy cone metric spaces. For $(x_1, x_2), (y_1, y_2) \in X_1 \times X_2$, let*

$$M((x_1, x_2), (y_1, y_2), t) = M_1(x_1, y_1, t) * M_2(x_2, y_2, t).$$

Then M is a fuzzy cone metric on $X_1 \times X_2$.

Proof. FCM1. Since $M_1(x_1, y_1, t) > 0$ and $M_2(x_2, y_2, t) > 0$, this implies that

$$M_1(x_1, y_1, t) * M_2(x_2, y_2, t) > 0.$$

Therefore

$$M((x_1, x_2), (y_1, y_2), t) > 0.$$

FCM2. Suppose that for all $t \gg \theta$, $M((x_1, x_2), (y_1, y_2), t) = 1$. This implies that $x_1 = y_1$ and $x_2 = y_2$ for all $t \gg \theta$. Hence

$$M_1(x_1, y_1, t) = 1$$

and

$$M_2(x_2, y_2, t) = 1.$$

It follows that

$$M((x_1, x_2), (y_1, y_2), t) = 1.$$

Conversely, suppose that $M((x_1, x_2), (y_1, y_2), t) = 1$. This implies that

$$M_1(x_1, y_1, t) * M_2(x_2, y_2, t) = 1.$$

Since

$$0 < M_1(x_1, y_1, t) \leq 1$$

and

$$0 < M_2(x_2, y_2, t) \leq 1,$$

it follows that

$$M_1(x_1, y_1, t) = 1$$

and

$$M_2(x_2, y_2, t) = 1.$$

Thus $x_1 = y_1$ and $x_2 = y_2$. Therefore $(x_1, x_2) = (y_1, y_2)$.

FCM3. To prove that $M((x_1, x_2), (y_1, y_2), t) = M((y_1, y_2), (x_1, x_2), t)$ we observe that

$$M_1(x_1, y_1, t) = M_1(y_1, x_1, t)$$

and

$$M_2(x_2, y_2, t) = M_2(y_2, x_2, t).$$

It follows that for all $(x_1, x_2), (y_1, y_2) \in X_1 \times X_2$ and $t \gg \theta$

$$M((x_1, x_2), (y_1, y_2), t) = M((y_1, y_2), (x_1, x_2), t).$$

FCM4. Since $(X_1, M_1, *)$ and $(X_2, M_2, *)$ are fuzzy cone metric spaces, we have that

$$M_1(x_1, z_1, t + s) \geq M_1(x_1, y_1, t) * M_1(y_1, z_1, s)$$

and

$$M_2(x_2, z_2, t + s) \geq M_2(x_2, y_2, t) * M_2(y_2, z_2, s)$$

for all $(x_1, x_2), (y_1, y_2), (z_1, z_2) \in X_1 \times X_2$ and $t, s \gg \theta$. Therefore

$$\begin{aligned} M((x_1, x_2), (z_1, z_2), t + s) &= M_1(x_1, z_1, t + s) * M_2(x_2, z_2, t + s) \\ &\geq M_1(x_1, y_1, t) * M_1(y_1, z_1, s) * M_2(x_2, y_2, t) * M_2(y_2, z_2, s) \\ &\geq M_1(x_1, y_1, t) * M_2(x_2, y_2, t) * M_1(y_1, z_1, s) * M_2(y_2, z_2, s) \\ &\geq M((x_1, x_2), (y_1, y_2), t) * M((y_1, y_2), (z_1, z_2), s). \end{aligned}$$

FCM5. Note that $M_1(x_1, y_1, t)$ and $M_2(x_2, y_2, t)$ are continuous with respect to t and $*$ is continuous too. It follows that

$$M((x_1, x_2), (y_1, y_2), t) = M_1(x_1, y_1, t) * M_2(x_2, y_2, t)$$

is also continuous. □

Proposition 3.8. *Let $(X_1, M_1, *)$ and $(X_2, M_2, *)$ be fuzzy cone metric spaces. We define*

$$M((x_1, x_2), (y_1, y_2), t) = M_1(x_1, y_1, t) * M_2(x_2, y_2, t).$$

*Then M is a complete fuzzy cone metric on $X_1 \times X_2$ if and only if $(X_1, M_1, *)$ and $(X_2, M_2, *)$ are complete.*

Proof. Suppose that $(X_1, M_1, *)$ and $(X_2, M_2, *)$ are complete fuzzy cone metric spaces. Let $\{a_n\}$ be a Cauchy sequence in $X_1 \times X_2$. Note that

$$a_n = (x_1^n, x_2^n)$$

and

$$a_m = (x_1^m, x_2^m).$$

Also, $M(a_n, a_m, t)$ converges to 1. This implies that

$$M((x_1^n, x_2^n), (x_1^m, x_2^m), t)$$

converges to 1 for each $t \gg \theta$. It follows that

$$M_1(x_1^n, x_1^m, t) * M_2(x_2^n, x_2^m, t)$$

converges to 1 for each $t \gg \theta$. Thus $M_1(x_1^n, x_1^m, t)$ converges to 1 and also $M_2(x_2^n, x_2^m, t)$ converges to 1. Therefore $\{x_1^n\}$ is a Cauchy sequence in $(X_1, M_1, *)$ and $\{x_2^n\}$ is a Cauchy sequence in $(X_2, M_2, *)$. Since $(X_1, M_1, *)$ and $(X_2, M_2, *)$ are complete fuzzy cone metric spaces, there exist $x_1 \in X_1$ and $x_2 \in X_2$ such that $M_1(x_1^n, x_1, t)$ converges to 1 and $M_2(x_2^n, x_2, t)$ converges to 1 for each $t \gg \theta$. Let $a = (x_1, x_2)$. Then $a \in X_1 \times X_2$. It follows that $M(a_n, a, t)$ converges to 1 for each $t \gg \theta$. This shows that $(X_1 \times X_2, M, *)$ is complete.

Conversely, suppose that $(X_1 \times X_2, M, *)$ is complete. We shall show that $(X_1, M_1, *)$ and $(X_2, M_2, *)$ are complete. Let $\{x_1^n\}$ and $\{x_2^n\}$ be Cauchy sequences in $(X_1, M_1, *)$ and $(X_2, M_2, *)$ respectively. Thus $M_1(x_1^n, x_1^m, t)$ converges to 1 and $M_2(x_2^n, x_2^m, t)$ converges to 1 for each $t \gg \theta$. It follows that

$$M((x_1^n, x_2^n), (x_1^m, x_2^m), t) = M_1(x_1^n, x_1^m, t) * M_2(x_2^n, x_2^m, t)$$

converges to 1. Then (x_1^n, x_2^n) is a Cauchy sequence in $X_1 \times X_2$. Since $(X_1 \times X_2, M, *)$ is complete, there exists a pair $(x_1, x_2) \in X_1 \times X_2$ such that $M((x_1^n, x_2^n), (x_1, x_2), t)$ converges to 1. Clearly, $M_1(x_1^n, x_1, t)$ converges to 1 and $M_2(x_2^n, x_2, t)$ converges to 1. Hence $(X_1, M_1, *)$ and $(X_2, M_2, *)$ are complete. This completes the proof. \square

Theorem 3.9. *Every separable fuzzy cone metric space is second countable.*

Proof. Let $(X, M, *)$ be the given separable fuzzy cone metric space. Let $A = \{a_n : n \in \mathbb{N}\}$ be a countable dense subset of X . Consider

$$B = \left\{ B \left(a_j, \frac{1}{k}, \frac{t_1}{k \|t_1\|} \right) : j, k \in \mathbb{N} \right\}$$

where $t_1 \gg \theta$ is constant. Then B is countable. We claim that B is a base for the family of all open sets in X . Let G be an open set in X . Let $x \in G$; then there exists r with $0 < r < 1$ and $t \gg \theta$ such that $B(x, r, t) \subset G$. Since $r \in (0, 1)$, we can find an $s \in (0, 1)$ such that $(1 - s) * (1 - s) > (1 - r)$. Choose $m \in \mathbb{N}$ such that $1/m < s$ and $t_1/m \|t_1\| \ll \frac{t}{2}$. Since A is dense in X , there exists an $a_j \in A$ such that $a_j \in B(x, 1/m, t_1/m \|t_1\|)$. Now if $y \in B(a_j, 1/m, t_1/m \|t_1\|)$, then

$$\begin{aligned} M(x, y, t) &\geq M \left(x, a_j, \frac{t}{2} \right) * M \left(y, a_j, \frac{t}{2} \right) \\ &\geq M \left(x, a_j, \frac{t_1}{m \|t_1\|} \right) * M \left(y, a_j, \frac{t_1}{m \|t_1\|} \right) \\ &\geq \left(1 - \frac{1}{m} \right) * \left(1 - \frac{1}{m} \right) \\ &\geq (1 - s) * (1 - s) \\ &> (1 - r). \end{aligned}$$

Thus $y \in B(x, y, t)$ and hence B is a basis. \square

Proposition 3.10. *A subspace of a separable fuzzy cone metric space is separable.*

Proof. Let X be a separable fuzzy cone metric space and Y a subspace of X . Let $A = \{x_n : n \in \mathbb{N}\}$ be a countable dense subset of X . For arbitrary but fixed $n, k \in \mathbb{N}$, if there are points $x \in X$ such that $M(x_n, x, t_1/k \|t_1\|) > 1 - 1/k$, where $t_1 \gg \theta$ is constant, choose one of them and denote it by x_{nk} . Let $B = \{x_{nk} : n, k \in \mathbb{N}\}$; then B is countable. Now we claim that $Y \subset \overline{B}$. Let $y \in Y$. Given r with $0 < r < 1$ and $t \gg \theta$ we can find a $k \in \mathbb{N}$ such that $(1 - 1/k) * (1 - 1/k) > 1 - r$ and $t_1/k \|t_1\| \ll \frac{t}{2}$. Since A is dense

in X , there exists an $m \in \mathbb{N}$ such that $M(x_m, y, t_1/k \|t_1\|) > 1 - 1/k$. But by definition of B , there exists an x_{mk} such that $M(x_{mk}, x_m, t_1/k \|t_1\|) > 1 - 1/k$. Now

$$\begin{aligned} M(x_{mk}, y, t) &\geq M\left(x_{mk}, x_m, \frac{t}{2}\right) * M\left(x_m, y, \frac{t}{2}\right) \\ &\geq M\left(x_{mk}, x_m, \frac{t_1}{k \|t_1\|}\right) * M\left(x_m, y, \frac{t_1}{k \|t_1\|}\right) \\ &\geq \left(1 - \frac{1}{k}\right) * \left(1 - \frac{1}{k}\right) \\ &> 1 - r. \end{aligned}$$

Thus $y \in \overline{B}$ and hence Y is separable. □

References

- [1] T. Bag, *Fuzzy cone metric spaces and fixed point theorems of contractive mappings*, Ann. Fuzzy Math. Inform., **6** (2013), 657–668.1
- [2] Z. Deng, *Fuzzy pseudo-metric spaces*, J. Math. Anal. Appl., **86** (1982), 74–95.1
- [3] M. A. Erceg, *Metric spaces in fuzzy set theory*, J. Math. Anal. Appl., **69** (1979), 205–230.1
- [4] A. George, P. Veeramani, *On some results in fuzzy metric spaces*, Fuzzy Sets and Systems, **64** (1994), 395–399.1
- [5] A. George, P. Veeramani, *On some results of analysis for fuzzy metric spaces*, Fuzzy Sets and Systems, **90** (1997), 365–368.1
- [6] V. Gregori, S. Romaguera, *Some properties of fuzzy metric spaces*, Fuzzy Sets and Systems, **115** (2000), 485–489.1
- [7] L. G. Huang, X. Zhang, *Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl., **332** (2007), 1468–1476.2
- [8] O. Kaleva, S. Seikkala, *On fuzzy metric spaces*, Fuzzy Sets and Systems, **12** (1984), 215–229.1
- [9] O. Kramosil, J. Michalek, *Fuzzy metric and statistical metric spaces*, Kybernetika, **11** (1975), 336–344.1
- [10] T. Öner, M. B. Kandemir, B. Tanay, *Fuzzy cone metric spaces*, J. Nonlinear Sci. Appl., **8** (2015), 610–616.1, 2
- [11] S. Rezapour, R. Hambarani, *Some notes on the paper: "Cone metric spaces and fixed point theorems of contractive mappings"*, J. Math. Anal. Appl., **332** (2007), 1468–1476.2
- [12] B. Schweizer, A. Sklar, *Statistical metric spaces*, Pacific J. Math., **10** (1960), 313–334.2
- [13] L. A. Zadeh, *Fuzzy sets*, Information and Control, **8** (1965), 338–353.1